



An initiative of the ABIM Foundation

American Society of Clinical Oncology



American Society of Clinical Oncology

Five Things Physicians and Patients Should Question

The American Society of Clinical Oncology (ASCO) is a medical professional oncology society committed to conquering cancer through research, education, prevention and delivery of high-quality patient care. ASCO recognizes the importance of evidence-based cancer care and making wise choices in the diagnosis and management of patients with cancer. After careful consideration by experienced oncologists, ASCO highlights ten categories of tests, procedures and/or treatments whose common use and clinical value are not supported by available evidence. These test and treatment options should not be administered unless the physician and patient have carefully considered if their use is appropriate in the individual case. As an example, when a patient is enrolled in a clinical trial, these tests, treatments and procedures may be part of the trial protocol and therefore deemed necessary for the patient's participation in the trial.

These items are provided solely for informational purposes and are not intended to replace a medical professional's independent judgment or as a substitute for consultation with a medical professional. Patients with any specific questions about the items on this list or their individual situation should consult their health care provider. New evidence may emerge following the development of these items. ASCO is not responsible for any injury or damage arising out of or related to any use of these items or to any errors or omissions.

1

Don't use cancer-directed therapy for solid tumor patients with the following characteristics: low performance status (3 or 4), no benefit from prior evidence-based interventions, not eligible for a clinical trial, and no strong evidence supporting the clinical value of further anti-cancer treatment.

- Studies show that cancer directed treatments are likely to be ineffective for solid tumor patients who meet the above stated criteria.
- Exceptions include patients with functional limitations due to other conditions resulting in a low performance status or those with disease characteristics (e.g., mutations) that suggest a high likelihood of response to therapy.
- Implementation of this approach should be accompanied with appropriate palliative and supportive care.

2

Don't perform PET, CT, and radionuclide bone scans in the staging of early prostate cancer at low risk for metastasis.

- Imaging with PET, CT, or radionuclide bone scans can be useful in the staging of specific cancer types. However, these tests are often used in the staging evaluation of low-risk cancers, despite a lack of evidence suggesting they improve detection of metastatic disease or survival.
- Evidence does not support the use of these scans for staging of newly diagnosed low grade carcinoma of the prostate (Stage T1c/T2a, prostate-specific antigen (PSA) <10 ng/ml, Gleason score less than or equal to 6) with low risk of distant metastasis.
- Unnecessary imaging can lead to harm through unnecessary invasive procedures, over-treatment, unnecessary radiation exposure, and misdiagnosis.

3

Don't perform PET, CT, and radionuclide bone scans in the staging of early breast cancer at low risk for metastasis.

- Imaging with PET, CT, or radionuclide bone scans can be useful in the staging of specific cancer types. However, these tests are often used in the staging evaluation of low-risk cancers, despite a lack of evidence suggesting they improve detection of metastatic disease or survival.
- In breast cancer, for example, there is a lack of evidence demonstrating a benefit for the use of PET, CT, or radionuclide bone scans in asymptomatic individuals with newly identified ductal carcinoma in situ (DCIS), or clinical stage I or II disease.
- Unnecessary imaging can lead to harm through unnecessary invasive procedures, over-treatment, unnecessary radiation exposure, and misdiagnosis.

4

Don't perform surveillance testing (biomarkers) or imaging (PET, CT, and radionuclide bone scans) for asymptomatic individuals who have been treated for breast cancer with curative intent.

- Surveillance testing with serum tumor markers or imaging has been shown to have clinical value for certain cancers (e.g., colorectal). However for breast cancer that has been treated with curative intent, several studies have shown there is no benefit from routine imaging or serial measurement of serum tumor markers in asymptomatic patients.
- False-positive tests can lead to harm through unnecessary invasive procedures, over-treatment, unnecessary radiation exposure, and misdiagnosis.

5

Don't use white cell stimulating factors for primary prevention of febrile neutropenia for patients with less than 20 percent risk for this complication.

- ASCO guidelines recommend using white cell stimulating factors when the risk of febrile neutropenia, secondary to a recommended chemotherapy regimen, is approximately 20 percent and equally effective treatment programs that do not require white cell stimulating factors are unavailable.
- Exceptions should be made when using regimens that have a lower chance of causing febrile neutropenia if it is determined that the patient is at high risk for this complication (due to age, medical history, or disease characteristics).

Disclaimer: These items are provided solely for informational purposes and are not intended to replace a medical professional's independent judgement or as a substitute for consultation with a medical professional. Patients with any specific questions about the items on this list or their individual situation should consult their health care provider.

Five More Things Physicians and Patients Should Question

6

Don't give patients starting on a chemotherapy regimen that has a low or moderate risk of causing nausea and vomiting antiemetic drugs intended for use with a regimen that has a high risk of causing nausea and vomiting.

- Over the past several years, a large number of effective drugs with fewer side effects have been developed to prevent nausea and vomiting from chemotherapy. When successful, these medications can help patients avoid spending time in the hospital, improve their quality of life and lead to fewer changes in the chemotherapy regimen.
- Oncologists customarily use different antiemetic drugs depending on the likelihood (low, moderate or high) for a particular chemotherapy program to cause nausea and vomiting. For chemotherapy programs that are likely to produce severe and persistent nausea and vomiting, there are new agents that can prevent this side effect. However, these drugs are very expensive and not devoid of side effects. For this reason, these drugs should be used only when the chemotherapy drugs that have a high likelihood of causing severe or persistent nausea and vomiting.
- When using chemotherapy that is less likely to cause nausea and vomiting, there are other effective drugs available at a lower cost.

7

Don't use combination chemotherapy (multiple drugs) instead of chemotherapy with one drug when treating an individual for metastatic breast cancer unless the patient needs a rapid response to relieve tumor-related symptoms.

- Although chemotherapy with multiple drugs, or combination chemotherapy, for metastatic breast cancer may slow tumor growth for a somewhat longer time than occurs when treating with a single agent, use of combination chemotherapy has not been shown to increase overall survival. In fact, the trade-offs of more frequent and severe side effects may have a net effect of worsening a patient's quality of life, necessitating a reduction in the dose of chemotherapy.
- Combination chemotherapy may be useful and worth the risk of more side effects in situations in which the cancer burden must be reduced quickly because it is causing significant symptoms or is life threatening. As a general rule, however, giving effective drugs one at a time lowers the risk of side effects, may improve a patient's quality of life, and does not typically compromise overall survival.

8

Avoid using PET or PET-CT scanning as part of routine follow-up care to monitor for a cancer recurrence in asymptomatic patients who have finished initial treatment to eliminate the cancer unless there is high-level evidence that such imaging will change the outcome.

- PET and PET-CT are used to diagnose, stage and monitor how well treatment is working. Available evidence from clinical studies suggests that using these tests to monitor for recurrence does not improve outcomes and therefore generally is not recommended for this purpose.
- False positive tests can lead to unnecessary and invasive procedures, overtreatment, unnecessary radiation exposure and incorrect diagnoses.
- Until high level evidence demonstrates that routine surveillance with PET or PET-CT scans helps prolong life or promote well-being after treatment for a specific type of cancer, this practice should not be done.

9

Don't perform PSA testing for prostate cancer screening in men with no symptoms of the disease when they are expected to live less than 10 years.

- Since PSA levels in the blood have been linked with prostate cancer, many doctors have used repeated PSA tests in the hope of finding "early" prostate cancer in men with no symptoms of the disease. Unfortunately, PSA is not as useful for screening as many have hoped because many men with prostate cancer do not have high PSA levels, and other conditions that are not cancer (such as benign prostate hyperplasia) can also increase PSA levels.
- Research has shown that men who receive PSA testing are less likely to die specifically from prostate cancer. However when accounting for deaths from all causes, no lives are saved, meaning that men who receive PSA screening have not been shown to live longer than men who do not have PSA screening. Men with medical conditions that limit their life expectancy to less than 10 years are unlikely to benefit from PSA screening as their probability of dying from the underlying medical problem is greater than the chance of dying from asymptomatic prostate cancer.

10

Don't use a targeted therapy intended for use against a specific genetic aberration unless a patient's tumor cells have a specific biomarker that predicts an effective response to the targeted therapy.

- Unlike chemotherapy, targeted therapy can significantly benefit people with cancer because it can target specific gene products, i.e., proteins that cancer cells use to grow and spread, while causing little or no harm to healthy cells. Patients who are most likely to benefit from targeted therapy are those who have a specific biomarker in their tumor cells that indicates the presence or absence of a specific gene alteration that makes the tumor cells susceptible to the targeted agent.
- Compared to chemotherapy, the cost of targeted therapy is generally higher, as these treatments are newer, more expensive to produce and under patent protection. In addition, like all anti-cancer therapies, there are risks to using targeted agents when there is no evidence to support their use because of the potential for serious side effects or reduced efficacy compared with other treatment options.

Abbreviations

CT, computed tomography; DCIS, ductal carcinoma in situ; PET, positron emission tomography; PSA, prostate-specific antigen.

How This List Was Created (1–5)

The American Society of Clinical Oncology (ASCO) has had a standing Cost of Cancer Care Task Force since 2007. The role of the Task Force is to assess the magnitude of rising costs of cancer care and develop strategies to address these challenges. In response to the 2010 New England Journal of Medicine article by Howard Brody, MD, “Medicine’s Ethical Responsibility for Health Care Reform – the Top Five List,” a subcommittee of the Cost of Cancer Care Task Force began work to identify common practices in oncology that were both common as well as lacking sufficient evidence for widespread use. Upon joining the Choosing Wisely campaign, the members of the subcommittee conducted a literature search to ensure the proposed list of items were supported by available evidence in oncology; ultimately the proposed Top Five list was approved by the full Task Force. The initial draft list was then presented to the ASCO Clinical Practice Committee, a group composed of community-based oncologists as well as the presidents of the 48 state/regional oncology societies in the United States. Advocacy groups were also asked to weigh in to ensure the recommendations would achieve the dual purpose of increasing physician-patient communication and changing practice patterns. A plurality of more than 200 clinical oncologists reviewed, provided input and supported the list. The final Top Five list in oncology was then presented to, discussed and approved by the Executive Committee of the ASCO Board of Directors and published in the Journal of Clinical Oncology. ASCO’s disclosure and conflict of interest policies can be found at www.asco.org.

How This List Was Created (6–10)

To guide ASCO in developing this list, suggestions were elicited from current ASCO committee members (approximately 700 individuals); 115 suggestions were received. After removing duplicates, researching the literature and discussing practice patterns, the Value in Cancer Care Task Force culled the list to 11 items, which comprised an ASCO Top Five voting slate that was sent back to the membership of all standing committees. Approximately 140 oncologists from its leadership cadre voted, providing ASCO with an adequate sample size and perspective on what oncologists find to be of little value. The list was reviewed and finalized by the Value in Cancer Care Task Force and ultimately reviewed and approved by the ASCO Board of Directors and published in the *Journal of Clinical Oncology*. ASCO’s disclosure and conflict of interest policies can be found at www.asco.org.

Sources

- 1 Azzoli CG, Temin S, Aliff T, et al: 2011 focused update of 2009 American Society of Oncology clinical practice guideline update on chemotherapy for stage IV non–small cell lung cancer. *J Clin Oncol* 29:3825–3831, 2011
Ettinger DS, Akerley W, Bepler G, et al: Non-small cell lung cancer. *J Natl Compr Canc Netw* 8:740–801, 2010
Carlson RW, Allred DC, Anderson BO, et al: Breast cancer. *J Natl Compr Canc Netw* 7:122–192, 2009
Engstrom PF, Benson AB 3rd, Chen YJ, et al: Colon cancer clinical practice guidelines. *J Natl Compr Canc Netw* 3:468–491, 2005
Smith TJ, Hillner BE: Bending the cost curve in cancer care. *N Engl J Med* 364:2060–2065, 2011
Peppercorn JM, Smith TJ, Helft PR, et al: American Society of Clinical Oncology statement: Toward individualized care for patients with advanced cancer. *J Clin Oncol* 29:755–760, 2011
- 2 Makarov DV, Desai RA, Yu JB, et al: The population level prevalence and correlates of appropriate and inappropriate imaging to stage incident prostate cancer in the Medicare population. *J Urol* 187:97-102, 2012
National Comprehensive Cancer Network: NCCN clinical practice guidelines in oncology (NCCN guidelines)-Prostate cancer. Version 4.2011
Thompson I, Thrasher JB, Aus G, et al: Guideline for the management of clinically localized prostate cancer: 2007 update. *J Urol* 177:2106–2130, 2007
- 3 Carlson RW, Allred DC, Anderson BO, et al: Invasive breast cancer. *J Natl Compr Canc Netw* 9:136–222, 2011
- 4 Locker GY, Hamilton S, Harris J, et al: ASCO 2006 update of recommendations for the use of tumor markers in gastrointestinal cancer. *J Clin Oncol* 24:5313–5327, 2006
Desch CE, Benson AB 3rd, Somerfield MR, et al: Colorectal cancer surveillance: 2005 update of an American Society of Clinical Oncology practice guideline. *J Clin Oncol* 23:8512-8519, 2005
Carlson RW, Allred DC, Anderson BO, et al: Breast cancer. *J Natl Compr Canc Netw* 7:122–192, 2009
Khatcheressian JL, Wolff AC, Smith TJ, et al: American Society of Clinical Oncology 2006 update of the breast cancer follow-up and management guideline in the adjuvant setting. *J Clin Oncol* 24: 5091–5097, 2006
Harris L, Fritsche H, Mennel R, et al: American Society of Clinical Oncology 2007 update of recommendations for the use of tumor markers in breast cancer. *J Clin Oncol* 25:5287–5312, 2007
- 5 Smith TJ, Khatcheressian J, Lyman GH, et al: ASCO 2006 update of recommendations for the use of white blood cell growth factors: An evidence based clinical practice guideline. *J Clin Oncol* 24:3187–3205, 2006
- 6 Basch E, Prestrud AA, Hesketh PJ, Kris MG, Feyer PC, Somerfield MR, Chesney M, Clark-Snow RA, Flaherty AM, Freundlich B, Morrow G, Rao KV, Schwartz RN, Lyman GH; American Society of Clinical Oncology. Antiemetics: American Society of Clinical Oncology clinical practice guideline update. *J Clin Oncol*. 2011 Nov 1;29:4189–98.
Saito M, Aogi K, Sekine I, Yoshizawa H, Yanagita Y, Sakai H, Inoue K, Kitagawa C, Ogura T, Mitsuhashi S. Palonosetron plus dexamethasone versus granisetron plus dexamethasone for prevention of nausea and vomiting during chemotherapy: a double-blind, double-dummy, randomized, comparative phase III trial. *Lancet Oncol*. 2009 Feb;10(2):115–24.
Aapro M, Fabi A, Nole F, Medici M, Steger G, Bachmann C, Roncoroni S, Roila F. Double-blind, randomised, controlled study of the efficacy and tolerability of palonosetron plus dexamethasone for 1 day with or without dexamethasone on days 2 and 3 in the prevention of nausea and vomiting induced by moderately emetogenic chemotherapy. *Ann Oncol*. 2010 May;21(5):1083–8.
Yu Z, Liu W, Wang L, Liang H, Huang Y, Si X, Zhang H, Liu D, Zhang H. The efficacy and safety of palonosetron compared with granisetron in preventing highly emetogenic chemotherapy-induced vomiting in the Chinese cancer patients: a phase II, multicenter, randomized, double-blind, parallel, comparative clinical trial. *Support Care Cancer*. 2009 Jan;17(1):99–102.

7

Cardoso F, Costa A, Norton L, Cameron D, Cufer T, Fallowfield L, Francis P, Gligorov J, Kyriakides S, Lin N, Pagani O, Senkus E, Thomssen C, Aapro M, Bergh J, Di Leo A, El Saghir N, Ganz PA, Gelmon K, Goldhirsch A, Harbeck N, Houssami N, Hudis C, Kaufman B, Leadbeater M, Mayer M, Rodger A, Rugo H, Sacchini V, Sledge G, van't Veer L, Viale G, Krop I, Winer E. 1st International consensus guidelines for advanced breast cancer (ABC 1). *Breast*. 2012 Jun;21(3):242–52.

Carrick S, Parker S, Thornton CE, Ghersi D, Simes J, Wilken N. Single agent versus combination chemotherapy for metastatic breast cancer. *Cochrane Database Syst Rev*. 2009 Apr 15;(2):CD003372.

National Comprehensive Cancer Network: NCCN clinical practice guidelines in oncology (NCCN Guidelines); breast cancer version: 1.2013.

Slamon DJ, Leyland-Jones B, Shak S, Fuchs H, Paton V, Bajamonde A, Fleming T, Eiermann W, Wolter J, Pegram M, Baselga J, Norton L. Use of chemotherapy plus a monoclonal antibody against HER2 for metastatic breast cancer that overexpresses HER2. *N Engl J Med*. 2001 Mar 15;344(11):783–92.

Howell A, Robertson JF, Quaresima Albano J, Aschermannova A, Mauriac L, Kleeberg UR, Vergote I, Erikstein B, Webster A, Morris C. Fulvestrant, formerly ICI 182,780, is as effective as anastrozole in postmenopausal women with advanced breast cancer progressing after prior endocrine treatment. *J Clin Oncol*. 2002 Aug 15;20(16):3396–403.

Lutz S, Berk L, Chang E, Chow E, Hahn C, Hoskin P, Howell D, Konski A, Kachnic L, Lo S, Sahgal A, Silverman L, von Gunten C, Mendel E, Vassil A, Bruner DW, Hartsell W; American Society for Radiation Oncology (ASTRO). Palliative radiotherapy for bone metastases: an ASTRO evidence-based guideline. *Int J Radiat Oncol Biol Phys*. 2011 Mar 15;79(4):965–76.

8

Phurrough S, Cano C, Dei Cas R, Ballantine L, Carino T; Centers for Medicare and Medicaid Services. Decision memo for positron emission tomography (FDG) for solid tumors (CAG–00181R4). Baltimore (MD): Centers for Medicare and Medicaid Services; 2003 Jul 8. 55 p. Report No.: CAG–00106R.

PET imaging in Ontario [Internet]. Ontario (CA): Cancer Care Ontario; 2012 May 28 [cited 26 Sep 2013]. Available from: www.cancercare.on.ca/ocs/clinicalprogs/imaging/pet.

Labianca R, Nordlinger B, Beretta GD, Brouquet A, Cervantes A; ESMO Guidelines Working Group. Primary colon cancer: ESMO Clinical Practice Guidelines for diagnosis, adjuvant treatment and follow-up. *Ann Oncol*. 2010 May;21 Suppl 5:v70–v7.

9

Raghavan D. PSA – Please Stop Agonizing (over prostate-specific antigen interpretation). *Mayo Clin Proc*. 2013 Jan;88:1–3.

Schroder FH, Hugosson J, Roobol MJ, Tammela TL, Ciatto S, Nelen V, Kwiatkowski M, Lujan M, Lilja H, Zappa M, Denis LJ, Recker F, Páez A, Mänttinen L, Bangma CH, Aus G, Carlsson S, Villers A, Rebillaud X, van der Kwast T, Kujala PM, Blijenberg BG, Stenman UH, Huber A, Taari K, Hakama M, Moss SM, de Koning HJ, Auvinen A; ERSPC Investigators. Prostate-cancer mortality at 11 years of follow-up. *N Engl J Med*. 2012 Mar 15;366(11):981–90.

Hugosson J, Carlsson S, Aus G, Bergdahl S, Khatami A, Lodding P, Pihl C-G, Stranne J, Holmberg E, Lilja H. Mortality results from the Göteborg randomized population-based prostate-cancer screening trial. *Lancet Oncol*. 2010 Aug;11(8):725–32.

Andriole GL, Crawford ED, Grubb RL III, Buys SS, Chia D, Church TR, Fouad MN, Gelmann EP, Kvale PA, Reding DJ, Weissfeld JL, Yokochi LA, O'Brien B, Clapp JD, Rathmell JM, Riley TL, Hayes RB, Kramer BS, Izmirlian G, Miller AB, Pinsky PF, Prorok PC, Gohagan JK, Berg CD; PLCO Project Team. Mortality results from a randomized prostate-cancer screening trial. *N Engl J Med*. 2009 Mar 26;360(12):1310–9.

Moyer VA; U.S. Preventive Services Task Force. Screening for prostate cancer: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med*. 2012 Jul 17;157(2):1–15.

Qaseem A, Barry MJ, Denberg TD, Owens DK, Shekelle P; Clinical Guidelines Committee of the American College of Physicians. Screening for prostate cancer: A guidance statement from the Clinical Guidelines Committee of the American College of Physicians. *Ann Intern Med*. 2013 May 21;158(10):761–9.

Carter HB, Albertson PC, Barry MJ, Etzioni R, Freedland SJ, Greene KL, Holmberg L, Kantoff P, Konety BR, Murad MH, Penson DF, Zietman AL. Early detection of prostate cancer: AUA Guideline. *J Urol*. 2013 Aug;190(2):419–26.

Basch E, Oliver TK, Vickers A, Thompson I, Kantoff P, Parnes H, Loblaw DA, Roth B, Williams J, Nam RK. Screening for prostate cancer with prostate-specific antigen testing: American Society of Clinical Oncology provisional clinical opinion. *J Clin Oncol*. 2012 Aug 20;30(24):3020–5.

10

Shaw A, Kim D, Nakagawa K, Seto T, Crinó L, Ahn MJ, De Pas T, Besse B, Solomon BJ, Blackhall F, Wu YL, Thomas M, O'Byrne KJ, Moro-Sibilot D, Camidge DR, Mok T, Hirsh V, Riey GJ, Iyer S, Tassell V, Polli A, Wilner KD, Jänne PA. Crizotinib versus chemotherapy in advanced ALK-positive lung cancer. *N Engl J Med*. 2013 Jun 20;368(25):2385–94.

Sequist L, Yang J, Yamamoto N, O'Byrne K, Hirsh V, Mok T, Geater SL, Orlov S, Tsai CM, Boyer M, Su WC, Bennis J, Kato T, Gorbunova V, Lee KH, Shah R, Massey D, Zazulina V, Shahidi M, Schuler M. Phase III study of afatinib or cisplatin plus pemetrexed in patients with metastatic lung adenocarcinoma with EGFR mutations. *J Clin Oncol*. 2013 Sep 20;31(27):3327–3334.

Chapman P, Hauschild A, Robert C, Haanen JB, Ascierto P, Larkin J, Dummer R, Garbe C, Testori A, Maio M, Hogg D, Lorigan P, Lebbe C, Jouary T, Schadendorf D, Ribas A, O'Day SJ, Sosman JA, Kirkwood JM, Eggermont AM, Dreno B, Nolop K, Li J, Nelson B, Hou J, Lee RJ, Flaherty KT, McArthur GA; BRIM-3 Study Group. Improved survival with vemurafenib in melanoma with BRAF V600E mutation. *N Engl J Med*. 2011 Jun 30;364(26):2507–16.

Lynch T, Bell D, Sordella R, Gurubhagavatula S, Okimoto RA, Brannigan BW, Harris PL, Haserlat SM, Supko JG, Haluska FG, Louis DN, Christiani DC, Settleman J, Haber DA. Activating mutations in the epidermal growth factor receptor underlying responsiveness of non-small-cell lung cancer to gefitinib. *N Engl J Med*. 2004 May 20;350(21):2129–39.

Keedy V, Temin S, Somerfield M, Beasley MB, Johnson DH, McShane LM, Milton DT, Strawn JR, Wakelee HA, Giaccone G. American Society of Clinical Oncology provisional clinical opinion: epidermal growth factor receptor (EGFR) mutation testing for patients with advanced non-small-cell lung cancer considering first-line EGFR tyrosine kinase inhibitor therapy. *J Clin Oncol*. 2011 May 20;29(15):2121–7.

Allegra C, Jessup J, Somerfield M, Hamilton SR, Hammond EH, Hayes DF, McAllister PK, Morton RF, Schilsky RL. American Society of Clinical Oncology provisional clinical opinion: testing for KRAS gene mutations in patients with metastatic colorectal carcinoma to predict response to anti-epidermal growth factor receptor monoclonal antibody therapy. *J Clin Oncol*. 2009 Apr 20;27(12):2091–6.

About the ABIM Foundation

The mission of the ABIM Foundation is to advance medical professionalism to improve the health care system. We achieve this by collaborating with physicians and physician leaders, medical trainees, health care delivery systems, payers, policymakers, consumer organizations and patients to foster a shared understanding of professionalism and how they can adopt the tenets of professionalism in practice.

To learn more about the ABIM Foundation, visit www.abimfoundation.org.



About the American Society of Clinical Oncology

The American Society of Clinical Oncology (ASCO) is the world's leading professional organization representing physicians who care for people with cancer. With more than 30,000 members, ASCO is committed to improving cancer care through scientific meetings, educational programs and peer-reviewed journals. ASCO is supported by its affiliate organization, the Conquer Cancer Foundation, which funds ground-breaking research and programs that make a tangible difference in the lives of people with cancer. ASCO's membership is comprised of clinical oncologists from all oncology disciplines and sub-specialties including medical oncology, therapeutic radiology, surgical oncology, pediatric oncology, gynecologic oncology, urologic oncology, and hematology; physicians and health care professionals participating in approved oncology training programs; oncology nurses; and other health care practitioners with a predominant interest in oncology.

For more information, please visit www.asco.org.

