

Supervised Consumption Facilities — Review of the Evidence

Sharon Larson, PhD; Norma Padrón, PhD; Tyler Bogaczyk; Jennifer Mason

Main Line Health Center for Population Health Research at Lankenau Institute for Medical Research
and Jefferson College of Population Health



Background — Philadelphia, PA

- Philadelphia Department of Public Health (PDPH) reported 907 deaths attributed to overdose in 2016.¹
 - 46.8 overdose deaths per 100,000 city residents
 - 729 of the 907 attributed to opioids
 - More than 3x the rate of Chicago and 4x the rate of New York City
- PDPH estimates around 469,000 individuals used prescription opioids in the past year, and 168,000 individuals currently use prescription opioids.¹
- Findings from the 2017 Mayor’s Task Force report²:
 - Approximately 50,000 people in Philadelphia misused prescription pain medications in the past year.
 - Philadelphia has an estimated 70,000 current heroin users.

What are SCFs?

- A Supervised Consumption Facility (SCF) provides a designated space for drug use under the supervision of medical professionals.
- SCFs draw in populations who inject drugs intravenously, providing a touchpoint to engage with populations generally disconnected from other social and public health access points.
 - A SCF in Sydney, Australia, showed that 15,054 people registered to use the Medically Supervised Injection Center, and 10,538 of those had never accessed any local health service prior.³
- Today there are over 100 SCFs in 66 cities and 11 countries worldwide.⁴
 - There are currently no “sanctioned” SCFs in the United States.

Harm Reduction and SCFs

Overdose Mortality Reduction

- Evidence shows that SCFs have a substantial effect on reducing mortality from overdose.⁵
- A study on overdose mortality near the Insite facility in Vancouver, Canada, found a 35% reduction in mortality within 500 meters of the facility within 3 years of its opening.⁶
- Significant reductions in overdose mortality have been noted in European SCFs.
 - Reductions in mortality are not only sustained but increase over time.
 - Spain’s SCF has been credited with reducing overdose deaths by over 50%, from 1,833 in 1991 to 773 by 2008.⁷

Injection Cessation

- European research on SCF clients (i.e., people who inject drugs [PWID]) has shown increased understanding regarding hygienic and safe injecting practices, as well as a reduction in syringe sharing.^{8,9}
- Evidence from Vancouver’s Insite SCF has shown clients who regularly visit the facility and have contact with counselors were more likely to seek entry into addiction treatment services.
 - Insite participants (2003 - 2005) who were part of the Scientific Evaluation of the Supervised Injecting cohort had a 30% increase in detoxification service use referrals attributed to the SCF opening.¹⁰

Reduction of Infections (HIV, HCV)

- Bacterial infections remain a significant issue for PWIDs due to shared needles, old needles and haste in injecting to avoid detection.
- SCFs worldwide report reduced bacterial infection by providing clean injection equipment, cleaning wounds and identifying serious infections early.^{6,11}
- The Insite clinic has provided evidence that SCFs can reduce blood-borne disease transmission by providing clean needles and safer injection education.
 - Analysis of a closed, unsanctioned SCF in Vancouver, Canada, showed that the facility prevented around 30 HIV and 81 HCV cases among PWID annually.¹²
- Conservative estimates on the reduction of HCV and HIV cases for a hypothetical SCF in Montreal, Canada, demonstrated each additional SCF would prevent 11 cases of HIV and 65 cases of HCV annually.¹³


Automobile Crashes

- Drug-impaired driving is an increasingly serious issue. In 2009, 32.8% of fatally injured drivers tested positive for drugs in the United States, and this rate increased in 2015 to 43%.¹⁴
- Automobile crashes caused by drivers who overdose on opioids are becoming so common that rescue crews in some areas are immediately administering naloxone to unconscious drivers.¹⁵
- However, we found no evidence reported in the available literature to support or negate a hypothesis that there is increased or decreased drugged driving in communities with a SCF.

Neighborhood Disorder

- Within 10 blocks of the Insite facility, Wood et al. (2004) found that the 12-week period after the facility’s opening was independently associated with reductions in the following numbers¹⁶:
 - injection drug users seen in public, from a daily mean of 4.3 to 2.4
 - publicly discarded syringes found, from a daily mean of 11.5 to 5.4
 - pieces of injection-related litter found, from a daily mean of 601 to 310

Incidence of Overdose Deaths in PHL by ZIP, 2016



Largest concentration was in the Kensington neighborhood.

Source: Medical Examiner’s Office, Philadelphia Department of Public Health

Potential Impacts of a SCF in Philadelphia

Variable	Low Case	High Case	Units
Population Health			
Estimated SCF-averted HIV infections	1	18	Cases
Estimated SCF-averted HCV infections	15	213	Cases
Marshall et al. (2011) estimated number of annual overdose deaths averted within 500m of SCF ⁶	27	48	Deaths
Milloy et al. (2008) estimated number of averted overdose deaths from opening a SCF ¹⁷	24	76	Deaths
Financial Impact			
Estimated annual savings due to SCF skin and soft tissue infection reduction	\$1,512,356	\$1,868,205	
Estimated total value of overdose deaths averted	\$12,462,213	\$74,773,276	
Estimated annual savings due to SCF reducing ambulance calls for overdose	\$123,776		
Estimated annual savings from keeping PWID out of emergency rooms	\$280,683		
Estimated annual savings on hospitals for PWID who overdose	\$247,971		

Conclusions

- SCFs provide cost savings, particularly in hospitalizations and other health services.
- SCFs prevent overdose deaths and reduce neighborhood disorder.
- The City of Philadelphia should develop a thoughtful evaluation that examines the implementation of these services.
 - Evaluation should pay attention to anticipated measurable outcomes that are directly or indirectly related to the selected interventions.

References

- PDPH, The Epidemic of Overdoses From Opioids in Philadelphia, in CHART. Philadelphia Department of Public Health. 2016.
- Philadelphia, C.o., The Mayor’s Task force to Combat the Opioid Epidemic in Philadelphia, Final Report and Recommendations. 2017: Philadelphia, PA.
- International Network of Drug Consumption Rooms. 2015; Available from: <http://www.drugconsumptionroom-international.org/>
- Rienzi, G. Supervised drug injection sites cut down on overdoses, infections, and opioid-related costs. *Johns Hopkins Magazine*. Fall 2017.
- Irwin, A., Jozaghi, E., Weir, B.W., Allen, S.T., Lindsay, A., & Sherman, S.G. Mitigating the heroin crisis in Baltimore, MD, USA: a cost-benefit analysis of a hypothetical supervised injection facility. *Harm Reduction Journal*, 2017;14(1), 29.
- Marshall, B.D., et al. Reduction in overdose mortality after the opening of North America’s first medically supervised safer injecting facility: a retrospective population-based study. *Lancet*. 2011; 377(9775): p. 1429-37.
- Ubelacker, S. Drug tests at B.C. supervised injection site found 80% contained fentanyl. The Canadian Press: Toronto. May, 2017.
- Peacy, J. Drug Consumption Rooms in Europe. Client Experience Survey in Amsterdam and Rotterdam. Regenboog Group, Correlation Network. 2014.
- Milloy, M.J. and Wood E. Emerging role of supervised injecting facilities in human immunodeficiency virus prevention. *Addiction*, 2009. 104(4): p. 620-21.
- Wood, E., et al. Rate of detoxification service use and its impact among a cohort of supervised injecting facility users. *Addiction*, 2007; 102(6): p. 916-19.
- Wood, E., et al. Impact of a medically supervised safer injecting facility on drug dealing and other drug-related crime. *Substance Abuse Treatment, Prevention, and Policy*, 2006; 1(1): p. 13.
- Jozaghi, E. Exploring the role of an unsanctioned, supervised peer driven injection facility in reducing HIV and hepatitis C infections in people that require assistance during injection. *Health & Justice*, 2015; 3(1): p. 16.
- Jozaghi, E., Reid, A.A., and Andresen, M.A. A cost-benefit/cost-effectiveness analysis of proposed supervised injection facilities in Montreal, Canada. *Substance Abuse Treatment, Prevention, and Policy*, 2014; 8(1): p. 25.
- GHSA, Drug Impaired Driving: A guide for states; 2017 update.
- Mitch, S. and Welsh-Huggins, A. Overdoses on the Road: Drugged driving rises as a menace. AP News. Aug. 17, 2017.
- Wood, E., et al. Changes in public order after the opening of a medically supervised safer injecting facility for illicit injection drug users. *Canadian Medical Association Journal*, 2004; 171(7): p. 731-34.
- Milloy, M.J., et al. Estimated drug overdose deaths averted by North America’s first medically-supervised safer injection facility. *PLoS One*, 2008. 3(10): p. e3351.